

TORIC DEGENERATIONS OF SPHERICAL VARIETIES

VALERY ALEXEEV AND MICHEL BRION

ABSTRACT. We prove that any affine, resp. polarized projective, spherical variety admits a flat degeneration to an affine, resp. polarized projective, toric variety. Motivated by Mirror Symmetry, we give conditions for the limit toric variety to be a Gorenstein Fano, and provide many examples. We also provide an explanation for the limits as boundary points of the moduli space of stable pairs whose existence is predicted by the Minimal Model Program.

INTRODUCTION

Let G be a connected reductive group. We prove that any affine, resp. polarized projective, spherical G -variety admits a flat degeneration to an affine, resp. polarized projective, toric variety. This is obtained as a corollary of a more general result which applies to arbitrary, not necessarily spherical G -schemes. We construct several degenerations, labeled by reduced decompositions $\underline{w}_0$ of the longest element w_0 in the Weyl group of G .

A number of special cases of this result have been proved before. The case of flag and Schubert varieties was considered by many authors, beginning with the work of Gonciulea and Lakshmibai [GL96] for grassmanians and varieties of complete flags, see also [Stu96, GŽ96, Chi00, KM03].

Caldero's paper [Cal02] constructs several degenerations of the flag variety G/B that depend on the choice of the reduced decomposition $\underline{w}_0$. His paper contains an important ingredient which we use for the general case. The choice of $\underline{w}_0$ previously appeared in the study of degenerations of the Bott-Samelson resolutions of G/B by Grossberg and Karshon [GK94]. As tempting as this connection is, the two degenerations do not seem to be directly related, see the discussion at the end of Section 6.

Regarding the more general case of spherical varieties, Kaveh [Kav03] proves the existence of a toric degeneration in the case where G is the symplectic group SP_{2n} .

We were led to the subject by trying to understand mirrors of Calabi-Yau hypersurfaces in spherical varieties, following the model suggested by Givental and others [Giv97, BCFKvS00]. In the latter paper, the four authors argue that mirrors of Calabi-Yau hypersurfaces in a partial flag variety $X = \mathrm{SL}_n/P$ can be found among Calabi-Yau hypersurfaces in X_0^* , the toric variety mirror-dual to the Gonciulea-Lakshmibai toric degeneration X_0 of X .

Therefore, in the projective case we pay special attention to the question of when the toric limits are Gorenstein Fano varieties, i.e., when they correspond to reflexive polytopes. For an arbitrary polarized spherical variety (X, L) , with toric degeneration (X_0, L_0) , we describe the corresponding polytope $Q = Q_{\underline{w}_0}(X, L)$. An important feature of the general case is that L_0 may only be a $\mathbb{Q}$ -line bundle, even

Date: March 23, 2004.

if X is Gorenstein Fano and $L = \mathcal{O}(-K)$ is the anticanonical line bundle ; but we still have $L_0 = \mathcal{O}(-K)$ in this case, i.e., X_0 is $\mathbb{Q}$ -Gorenstein Fano.

In the case where $X = G/P$ is a flag variety and $L = \mathcal{O}(-K)$, we give a criterion for X_0 to be Gorenstein, i.e., for the polytope Q to be reflexive. In particular, if $X = \mathrm{SL}_n/P$ there are *many* different reflexive polytopes corresponding to toric degenerations and not just the Gelfand-Tsetlin polytope, which corresponds to the Gonciulea-Lakshmibai degeneration.

We conjecture that for groups of type A_n the limits of $(G/B, \mathcal{O}(-K))$ are Gorenstein. However, this fails for general types as we show in Example 5.11 for a simple group G of type E_6 .

The paper is organized as follows. Section 1 introduces the necessary technical tools: the dual canonical basis and its string parameterizations, string cones and polytopes, Caldero's filtration.

Section 2 is devoted to the affine case. We construct limits of an affine G -scheme X , which are toric if X is spherical. We also prove that these limits have rational singularities and trivial canonical class, if this holds for X .

In Section 3, we adapt these results to the setting of polarized projective varieties. In particular, we degenerate any polarized spherical variety (X, L) to a $\mathbb{Q}$ -polarized toric variety (X_0, L_0) , which is a $\mathbb{Q}$ -Gorenstein Fano if so is X . Further, we show that the moment polytope of (X_0, L_0) projects onto the moment polytope of (X, L) , with fibers being string polytopes. The latter are the moment polytopes of toric limits of flag varieties.

In Section 4 we describe these string and moment polytopes and give a criterion for their integrality, in particular, a criterion for the anticanonical limit of a flag variety to be a Gorenstein Fano variety. We prove that for the varieties G/P associated with minuscule or cominuscule weights, the limit toric varieties are Gorenstein Fano. In general, we show that the string polytopes have a number of special vertices where the associated toric variety is Gorenstein.

Section 5 presents examples and counterexamples of string polytopes, including the Gelfand-Tsetlin polytopes for classical groups. All of these are integral in type A , but not in other types. We also consider properties of string polytopes that do not depend on the choice of $\underline{w}_0$.

In Section 6, we explain how the above toric limits may be understood geometrically as boundary points of a “universal” compact moduli space of stable pairs, the existence of which is predicted by the Minimal Model Program.

Acknowledgment. The first author was partially supported by NSF grant No. 0101280.

1. NOTATION AND BACKGROUND

We begin by introducing notation that will be used throughout this paper. We consider algebraic schemes, varieties and groups over the field $\mathbb{C}$ of complex numbers. Let G be a connected reductive group, B a Borel subgroup with unipotent radical U , and T a maximal torus of B , so that $B = TU$. Let $\Phi = \Phi(G, T)$ be the root system of (G, T) , with subset of positive roots $\Phi^+ = \Phi(B, T)$. We denote by $\alpha_1, \dots, \alpha_r$ the corresponding simple roots, where r is the semisimple rank of G .

Let W be the Weyl group of (G, T) and let $s_1, \dots, s_r \in W$ be the simple reflections associated with $\alpha_1, \dots, \alpha_r$. These generators of W define the length function

ℓ on this group; let w_0 be the unique element of maximal length. Then $N := \ell(w_0)$ is the number of positive roots, i.e., the dimension of the flag variety G/B .

Denote by Λ the character group of T , also called the weight lattice of G . Let Λ^+ be the subset of dominant weights; it is stable under the involution

$$\lambda \mapsto -w_0\lambda =: \lambda^*$$

of Λ . There is a partial order on Λ , defined by $\mu \leq \lambda$ if there exist nonnegative integers $n_1, \dots, n_r$ such that $\lambda - \mu = n_1\alpha_1 + \dots + n_r\alpha_r$. We put $\Lambda_{\mathbb{R}} := \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$, then the convex cone generated by Λ^+ in $\Lambda_{\mathbb{R}}$ is the positive Weyl chamber $\Lambda_{\mathbb{R}}^+$. For $\lambda, \mu \in \Lambda^+$, we have $\mu \leq \lambda$ if and only if: μ lies in the convex hull of the Weyl group orbit $W\lambda$, and $\lambda - \mu$ lies in the root lattice. In particular, there are only finitely many $\mu \in \Lambda^+$ that are smaller than a fixed $\lambda \in \Lambda^+$.

Recall that Λ^+ parametrizes the isomorphism classes of simple (rational) G -modules, as follows. Consider the algebra

$$A := \mathbb{C}[G]^U$$

of regular functions on G , invariant under right multiplication by U . Then $G \times T$ acts on A , where G acts via left multiplication, and T acts on the right, since it normalizes U . We have an isomorphism of $G \times T$ -modules

$$A \cong \bigoplus_{\lambda \in \Lambda^+} V(\lambda^*),$$

where $V(\lambda^*)$ is a simple G -module of highest weight $\lambda^* = -w_0\lambda$, and T acts on each $V(\lambda^*)$ via the character λ . Thus, $V(\lambda^*)$ is the weight space A_{λ} .

The assignment $\lambda \mapsto V(\lambda^*)$ is the desired parametrization. Note that each $V(\lambda^*) \subset A$ has a canonical linear form, B -eigenvector of weight λ , obtained by restricting any function on G/U to $B/U \cong T$. In other words, $V(\lambda) = V(\lambda^*)^*$ has a canonical highest weight vector, denoted by v_{λ} . The stabilizer in G of the line spanned by v_{λ} is a parabolic subgroup containing B , denoted by P_{λ} . Then λ extends to a character of P_{λ} , that we still denote by λ ; let L_{λ} be the corresponding G -linearized line bundle on G/P_{λ} . We have

$$H^0(G/P_{\lambda}, L_{\lambda}) = V(\lambda^*).$$

Note that $P_{\lambda} = B$ if and only if λ is regular, i.e., in the interior of $\Lambda_{\mathbb{R}}^+$.

We next recall results from [Cal02]. The vector space A has a remarkable basis $\mathbb{B} = (b_{\lambda, \varphi})$, where each $b_{\lambda, \varphi}$ is an eigenvector of $T \times T \subseteq G \times T$, of weight λ for the right T -action. Specifically, for fixed λ , the vectors $b_{\lambda, \varphi}$ form the dual basis of the basis of $V(\lambda)$ consisting of the nonzero bv_{λ} , where b lies in the specialization at $q = 1$ of Kashiwara-Lusztig's *canonical basis* $\mathcal{B}$. We say that $\mathbb{B}$ is the *dual canonical basis* of A (its elements are denoted by $b_{\lambda, \varphi}^*$ in [Cal02], but we will drop the $*$ as we will only use the dual canonical basis). The parameters φ are defined via the *string parametrization* of $\mathbb{B}$ (or $\mathcal{B}$) which depends on the choice of a reduced decomposition

$$\underline{w}_0 = (s_{i_1}, s_{i_2}, \dots, s_{i_N})$$

of the longest element of W , i.e., $w_0 = s_{i_1} s_{i_2} \dots s_{i_N}$. It allows to define an injective map

$$\iota_{\underline{w}_0} : \mathbb{B} \rightarrow \Lambda^+ \times \mathbb{N}^N, \quad b_{\lambda, \varphi} \mapsto (\lambda, \varphi) = (\lambda, t_1, \dots, t_N).$$

The image of $\iota_{\underline{w}_0}$ is the intersection of a rational convex polyhedral cone $\mathcal{C}_{\underline{w}_0}$ of $\Lambda_{\mathbb{R}} \times \mathbb{R}^N$, with the lattice $\Lambda \times \mathbb{Z}^N$. The projection of $\mathcal{C}_{\underline{w}_0}$ to $\mathbb{R}^N$ is the *string cone* $C_{\underline{w}_0}$, a rational polyhedral convex cone of $\mathbb{R}^N$. Define a map

$$\pi : \Lambda_{\mathbb{R}} \times \mathbb{R}^N \rightarrow \mathbb{R}^N, \quad (\lambda, t_1, \dots, t_N) \mapsto -\lambda + t_1 \alpha_{i_1} + \dots + t_N \alpha_{i_N}.$$

Then the weight of $b_{\lambda, \varphi}$ as a left T -eigenvector equals $\pi(\lambda, t_1, \dots, t_N)$.

A crucial feature of the dual canonical basis is the following multiplicative property:

$$b_{\lambda, \varphi} b_{\mu, \psi} = b_{\lambda + \mu, \varphi + \psi} + \sum_{\gamma} c_{\lambda, \varphi; \mu, \psi}^{\gamma} b_{\lambda + \mu, \gamma},$$

where $c_{\lambda, \varphi; \mu, \psi}^{\gamma} = 0$ unless $\gamma < \varphi + \psi$. Here $\leq$ denotes the lexicographic ordering on $\mathbb{N}^N$. In particular, if a product $b_{\lambda, \varphi} b_{\mu, \psi}$ is a nonzero scalar multiple of a basis vector $b_{\nu, \gamma}$, then $\nu = \lambda + \mu$ and $\gamma = \varphi + \psi$.

From this, Caldero deduces the existence of an increasing filtration of the algebra A by $T \times T$ -submodules, such that the associated graded algebra, $\text{gr } A$, is isomorphic to the algebra of the monoid

$$\Gamma_{\underline{w}_0} := i_{\underline{w}_0}(\mathbb{B}) = \mathcal{C}_{\underline{w}_0} \cap (\Lambda \times \mathbb{Z}^N).$$

In geometric terms, the affine variety

$$G//U := \text{Spec}(A)$$

degenerates to

$$(G//U)_0 := \text{Spec } \mathbb{C}[\Gamma_{\underline{w}_0}].$$

The latter is an affine toric variety for the action of the torus $T \times \mathbb{T}$, where we put

$$\mathbb{T} := (\mathbb{C}^*)^N.$$

Further, the degeneration is compatible with the actions of $T \times T$ on $G//U$ (regarding $T \times T$ as a subgroup of $G \times T$), and on $(G//U)_0$ via the homomorphism of tori

$$T \times T \rightarrow T \times \mathbb{T}, \quad (x, y) \mapsto (x^{-1}y, \alpha_{i_1}(x), \dots, \alpha_{i_N}(x)).$$

Indeed, the induced homomorphism of character groups maps any $(\lambda, t_1, \dots, t_N) \in \Lambda \times \mathbb{Z}^N$ to $(-\lambda + t_1 \alpha_{i_1} + \dots + t_N \alpha_{i_N}, \lambda) \in \Lambda \times \Lambda$.

Actually, these results are obtained in [Cal02] for a simply-connected, semisimple group G , but their extension to an arbitrary connected reductive group G is immediate. Indeed, we have $G = (G^{\text{ss}} \times C)/Z$ where G^{ss} is a simply-connected, semisimple group, C is a torus, and Z is a finite central subgroup of $G^{\text{ss}} \times C$. Thus, $U \subset G^{\text{ss}}$, the Weyl groups of G and G^{ss} are identified, and $\mathbb{C}[G]^U$ is the subalgebra of Z -invariants in $\mathbb{C}[G^{\text{ss}}]^U \otimes \mathbb{C}[C]$. The latter space has a basis consisting of the $b_{\lambda, \varphi} \otimes \chi$, where $b_{\lambda, \varphi}$ lies in the dual canonical basis for G^{ss} , and χ is a character of C . Thus, a basis of $\mathbb{C}[G]^U$ consists of those $b_{\lambda, \varphi} \otimes \chi$ such that $\lambda + \chi$ restricts to the trivial character of Z .

The cone $\mathcal{C}_{\underline{w}_0}$ is described in [Lit98, §1] and [BZ01, 3.10]:

Theorem 1.1. *The cone $\mathcal{C}_{\underline{w}_0} \subset \Lambda_{\mathbb{R}} \times \mathbb{R}^N$ is the intersection of the preimage $\Lambda_{\mathbb{R}} \times C_{\underline{w}_0}$ of the string cone $C_{\underline{w}_0} \subset \mathbb{R}^N$ with the N half-spaces*

$$t_k \leq \langle \lambda, \alpha_{i_k}^{\vee} \rangle - \sum_{\ell=k+1}^N \langle \alpha_{i_\ell}, \alpha_{i_k}^{\vee} \rangle t_\ell, \quad k = 1, \dots, N.$$

Therefore, for every dominant weight λ the basis $(b_{\lambda, \varphi})$ is in bijection with the set of integral points of the rational convex polytope $Q_{\underline{w}_0}(\lambda)$ obtained by intersecting the string cone with the cone defined by the above λ -inequalities; here and later, by integral points we mean those in the lattice $\mathbb{Z}^N$. Note that we may define $Q_{\underline{w}_0}(\lambda)$, more generally, for any $\lambda \in \Lambda_{\mathbb{R}}^+$.

Definition 1.2. We will call the polytopes $Q_{\underline{w}_0}(\lambda)$ the *string polytopes*.

Let $\lambda \in \Lambda^+$. Under the projection

$$\pi_{\lambda} : \mathbb{R}^N \rightarrow \Lambda_{\mathbb{R}}, \quad (t_1, \dots, t_N) \mapsto -\lambda + t_1 \alpha_{i_1} + \dots + t_N \alpha_{i_N},$$

the integral points of $Q_{\underline{w}_0}(\lambda)$ map onto the weights of the T -module $V(\lambda^*)$. These weights are precisely those points μ in the convex hull of the orbit $W\lambda^* = -W\lambda$, such that $\lambda^* - \mu$ is in the root lattice. Applying this construction to all positive integral multiples $n\lambda$ and using the equality $Q_{\underline{w}_0}(n\lambda) = nQ_{\underline{w}_0}(\lambda)$, we see that in fact

$$\pi_{\lambda}(Q_{\underline{w}_0}(\lambda)) = \text{Conv}(W\lambda^*).$$

Moreover, for each $\mu \in W\lambda^*$, the μ -weight subspace of $V(\lambda^*)$ is 1-dimensional, so the point μ has exactly one preimage in $Q_{\underline{w}_0}(\lambda)$. This preimage must be a vertex, since μ is a vertex of $\text{Conv}(W\lambda^*)$.

Definition 1.3. The preimages $\pi_{\lambda}^{-1}(\mu)$, $\mu \in W\lambda^*$, will be called the *extremal weight vertices* of $Q_{\underline{w}_0}(\lambda)$ and denoted by q_{μ} .

Note that each q_{μ} is integral. For example, the *lowest weight vertex* $q_{-\lambda}$ is just the origin. If λ is regular, then the λ -inequalities are strict inequalities at the origin, so that the tangent cone to the string polytope at $q_{-\lambda}$ is the whole string cone.

On the other hand, the *highest weight vertex* q_{λ^*} is the point where all the λ -inequalities are equalities. One easily checks that

$$q_{\lambda^*} = (\langle \lambda^*, \beta_1^{\vee} \rangle, \dots, \langle \lambda^*, \beta_N^{\vee} \rangle),$$

where

$$\beta_1 := \alpha_{i_1}, \beta_2 := s_{i_1} \alpha_{i_2}, \dots, \beta_N := s_{i_1} \cdots s_{i_{N-1}} \alpha_{i_N}$$

is the enumeration of the simple roots associated with the reduced decomposition $\underline{w}_0$.

Explicit inequalities for the string cone $C_{\underline{w}_0}$ (and therefore for the string polytopes) in the case of groups of type A_n are given by Berenstein-Zelevinsky in [BZ01, Thm.3.14]. Littelmann [Lit98, Thm.4.2] describes the cone $C_{\underline{w}_0}$ for the so called *nice decompositions* $\underline{w}_0$, and this is generalized in [BZ01, Thm.3.12]. Also, [Lit98, §9] contains formulas for a few particular decompositions in types F_4 and E_8 .

By using [Lit98, Thm.4.2], one checks that the highest weight vertex is in the interior of the string cone, whenever λ is regular and $\underline{w}_0$ is a nice decomposition. Equivalently, the tangent cone to the string polytope at q_{λ^*} is defined by the inequalities

$$t_k + \sum_{\ell=k+1}^N \langle \alpha_{i_{\ell}}, \alpha_{i_k}^{\vee} \rangle t_{\ell} \leq 0, \quad k = 1, \dots, N.$$

In particular, this cone is generated by a basis of the lattice $\mathbb{Z}^N$. We do not know if this extends to arbitrary reduced decompositions and regular dominant weights.

2. DEGENERATIONS OF AFFINE G -ALGEBRAS

Let R be an affine G -algebra, i.e., a finitely generated algebra over $\mathbb{C}$ on which G acts rationally by algebra automorphisms. We first recall results on the structure of the G -module R , for which a general reference is [Gro97]. Let R^U be the subalgebra of U -invariants in R ; this is an affine T -algebra. Likewise, the algebra R^G of G -invariants is finitely generated, and each weight space R_λ^U is a finitely generated R^G -module; further, $R_0^U = R^B = R^G$. We have canonical isomorphisms of R^G - G -modules

$$R \cong \bigoplus_{\lambda \in \Lambda^+} \operatorname{Hom}^G(V(\lambda), R) \otimes V(\lambda) \cong \bigoplus_{\lambda \in \Lambda^+} R_\lambda^U \otimes V(\lambda),$$

that map any $u \otimes v \in \operatorname{Hom}^G(V(\lambda), R) \otimes V(\lambda)$ to $u(v) \in R$, resp. $u(v_\lambda) \otimes v \in R_\lambda^U \otimes V(\lambda)$. The corresponding summands of R are the *isotypical components* $R_{(\lambda)}$. Their products in R satisfy

$$R_{(\lambda)} R_{(\mu)} \subseteq \bigoplus_{\nu \in \Lambda^+, \nu \leq \lambda + \mu} R_{(\nu)}.$$

We now extend Caldero's filtration to this setting. Choose a reduced decomposition $\underline{w}_0$, then we obtain an isomorphism of R^G - T -modules

$$R \cong \bigoplus_{(\lambda, \varphi) \in \mathcal{C}_{\underline{w}_0}} R_\lambda^U \otimes b_{\lambda^*, \varphi},$$

where T acts on each space $R_\lambda^U \otimes b_{\lambda^*, t_1, \dots, t_N}$ via the weight $\pi_\lambda(t_1, \dots, t_N)$.

Define a partial ordering $\leq$ on $\Gamma_{\underline{w}_0}$ by setting: $(\mu, \psi) \leq (\lambda, \varphi)$ if, either $\mu < \lambda$, or $\mu = \lambda$ and $\psi \leq \varphi$. Note that this ordering is compatible with addition, and that there are only finitely many $(\mu, \psi) \in \Gamma_{\underline{w}_0}$ that are smaller than a given (λ, φ) . Now put

$$R_{\leq(\lambda, \varphi)} := \bigoplus_{(\mu, \psi) \leq (\lambda, \varphi)} R_\mu^U \otimes b_{\mu^*, \psi}$$

and define similarly $R_{<(\lambda, \varphi)}$. Both are finitely generated R^G -submodules of R , stable under T . Further, the $R_{\leq(\lambda, \varphi)}$ form an increasing, exhaustive filtration of R , satisfying the multiplicative properties

$$R_{\leq(\lambda, \varphi)} R_{\leq(\mu, \psi)} \subseteq R_{\leq(\lambda + \mu, \varphi + \psi)},$$

$$R_{\leq(\lambda, \varphi)} R_{<(\mu, \psi)} \subseteq R_{<(\lambda + \mu, \varphi + \psi)}.$$

Indeed, we have

$$(f_\lambda \otimes b_{\lambda^*, \varphi})(f_\mu \otimes b_{\mu^*, \psi}) \in (f_\lambda f_\mu \otimes b_{\lambda^* + \mu^*, \varphi + \psi}) + R_{<(\lambda + \mu, \varphi + \psi)},$$

as follows from the multiplicative properties of the isotypical components and of the dual canonical basis. Thus, we may define the associated graded algebra

$$\operatorname{gr} R := \bigoplus_{(\lambda, \varphi) \in \Gamma_{\underline{w}_0}} R_{\leq(\lambda, \varphi)} / R_{<(\lambda, \varphi)}.$$

This is a R^G - T -algebra, graded by the monoid $\Gamma_{\underline{w}_0}$; it may be regarded as the space $\bigoplus_{(\lambda, \varphi) \in \Gamma_{\underline{w}_0}} R_\lambda^U \otimes b_{\lambda^*, \varphi}$ with “componentwise” multiplication. This implies readily

Proposition 2.1. *With the preceding notation, the R^G - T -algebra $\text{gr } R$ is isomorphic to $(R^U \otimes \text{gr } A)^T$, where T acts on R^U naturally, and on $\text{gr } A$ via the inverse of its right action. As a consequence, $\text{gr } R$ is an affine $T \times \mathbb{T}$ -algebra, and its $\mathbb{T}$ -invariant subring is isomorphic to R^G .*

We would like to obtain $\text{gr } R$ as the associated graded algebra of R for a $\mathbb{N}$ -filtration, instead of the multifiltration indexed by the partially ordered monoid $\Gamma_{\underline{w}_0}$. This is the content of the following

Proposition 2.2. *There exist an affine $\mathbb{N}$ -graded T -algebra $\mathcal{R}$ and a T -invariant element $t \in \mathcal{R}_1$ such that*

- (i) *t is a nonzerodivisor in $\mathcal{R}$, i.e., $\mathcal{R}$ is flat over the polynomial ring $\mathbb{C}[t]$.*
- (ii) *The $\mathbb{C}[t, t^{-1}]$ - T -algebra $\mathcal{R}[t^{-1}]$ is isomorphic to $R[t, t^{-1}]$.*
- (iii) *The T -algebra $\mathcal{R}/t\mathcal{R}$ is isomorphic to $\text{gr } R$.*

Proof. We adapt the arguments of [Cal02, 3.2] to our setting. By Proposition 2.1, the algebra $\text{gr } R$ is finitely generated. Choose homogeneous generators $\bar{f}_1, \dots, \bar{f}_n$ of respective degrees $(\lambda_1, \varphi_1), \dots, (\lambda_n, \varphi_n)$, and lift these generators to $f_1, \dots, f_n \in R$. Then one easily checks that $f_1, \dots, f_n$ generate the algebra R . Let S denote the $\Gamma_{\underline{w}_0}$ -graded polynomial ring $\mathbb{C}[x_1, \dots, x_n]$, where $\deg(x_i) := (\lambda_i, \varphi_i)$. Choose homogeneous generators $\bar{g}_1, \dots, \bar{g}_p$ of the kernel of the surjective homomorphism $S \rightarrow \text{gr } R$, $x_i \mapsto \bar{f}_i$, and put $\deg(\bar{g}_i) = (\mu_i, \psi_i)$. Then $\bar{g}_i(f_1, \dots, f_n) \in R_{<(\mu_i, \psi_i)}$ for all i . Thus, we may find elements $g_1, \dots, g_p$ of S such that $g_i(f_1, \dots, f_n) = 0$ and $g_i \in \bar{g}_i + S_{<(\mu_i, \psi_i)}$. It follows that the natural map $S/g_1S + \dots + g_pS \rightarrow R$ is an isomorphism (since both sides are filtered algebras, and the associated graded map is an isomorphism).

Next consider all the differences $(\mu_i, \psi_i) - (\nu_j, \gamma_j)$, where (ν_j, γ_j) is the degree of a homogeneous component of $g_i - \bar{g}_i$. We claim that there exists a linear form $e : \Lambda_{\mathbb{R}} \times \mathbb{R}^N \rightarrow \mathbb{R}$ that takes positive integral values at all nonzero points of $\Lambda^+ \times \mathbb{N}^N$, and at all these differences. Indeed, recall that either $\mu_i > \nu_j$, or $\mu_i = \nu_j$ and $\psi_i > \gamma_j$. So we may take $e = (Ae_1, e_2)$, where A is a large positive integer, $e_1 : \Lambda_{\mathbb{R}} \rightarrow \mathbb{R}$ is a linear form that takes positive integral values at all positive roots, and $e_2 : \mathbb{R}^N \rightarrow \mathbb{R}$ is a linear form that takes positive integer values at all nonzero points of $\mathbb{N}^N$ and at all the (finitely many) points $\psi_i - \gamma_j$ such that $\lambda_i = \mu_j$ (the existence of e_2 follows from [Cal02, Lemma 3.2] and its proof).

Now put for any nonnegative integer m :

$$R_{\leq m} := \langle f_1^{a_1} \cdots f_n^{a_n} \mid e(\sum_{i=1}^n a_i(\mu_i, \psi_i)) \leq m \rangle \subset R.$$

This defines a T -stable $\mathbb{N}$ -filtration of the algebra R , such that the associated graded is our original $\text{gr } R$. Let t be an indeterminate and consider the Rees algebra

$$\mathcal{R} := \bigoplus_{m=0}^{\infty} R_{\leq m} t^m \subset R[t].$$

Then $\mathcal{R}$ and t have the desired properties; the finite generation of $\mathcal{R}$ follows from that of $\text{gr } R$. $\square$

(One easily checks that the Rees algebra associated with our multifiltration of R is not finitely generated. The reason is that the set of nonnegative elements of $\Lambda \times \mathbb{Z}^N$ spans a convex cone which is not polyhedral; it is not even closed.)

Proposition 2.2 may be formulated in geometric terms: the affine G -algebra R corresponds to an affine G -scheme $X = \operatorname{Spec}(R)$, and $\operatorname{gr} R$, to an affine $T \times \mathbb{T}$ -scheme that we denote by X_0 . Now there exists a family of affine T -schemes

$$\pi : \mathcal{X} \rightarrow \mathbb{A}^1$$

such that

- (i) π is flat.
- (ii) π is trivial with fiber X over the complement of 0 in $\mathbb{A}^1$.
- (iii) The fiber of π at 0 is isomorphic to X_0 .

We then say that X *degenerates to* X_0 , and that X_0 *is a limit of* X . Note that X_0 depends on the choice of a reduced decomposition of w_0 .

Next we consider an affine *spherical* G -variety $X = \operatorname{Spec}(R)$, i.e., X is normal and contains a dense B -orbit. Equivalently, R is a domain, and we have an isomorphism of G -modules

$$R \cong \bigoplus_{\lambda \in \Lambda^+ \cap \operatorname{Cone}(X)} V(\lambda),$$

where $\operatorname{Cone}(X)$ is a rational polyhedral convex cone in $\Lambda_{\mathbb{R}}^+$. In particular, the multiplicity of each $V(\lambda)$ in R is 0 or 1, so that R is *multiplicity-free*. Then $\operatorname{Cone}(X)$ is uniquely determined by X ; it is called the *weight cone* of X . Now Propositions 2.1 and 2.2 imply readily

Proposition 2.3. *Let X be an affine spherical G -variety with weight cone $\operatorname{Cone}(X)$. Then X degenerates to the affine toric $T \times \mathbb{T}$ -variety X_0 such that*

$$\operatorname{Cone}(X_0) = (\operatorname{Cone}(X) \times \mathbb{R}^N) \cap \mathcal{C}_{\underline{w}_0}.$$

This degeneration is T -equivariant, where T acts on X_0 via the homomorphism $T \rightarrow T \times \mathbb{T}$, $x \mapsto (x^{-1}, \alpha_{i_1}(x), \dots, \alpha_{i_N}(x))$.

Returning to an arbitrary G -scheme X with limit X_0 , many geometric properties hold for X if and only if they hold for X_0 , as can be shown along the lines of [Pop87, §6] (see also [Gro97, §18]). We record three such properties that will be of use in the sequel.

For a normal variety X , we denote by $\mathcal{O}(K_X)$ its canonical sheaf, where K_X is a canonical divisor; then the rational equivalence class of K_X is uniquely defined and called the *canonical class* of X . Also, recall that X *has rational singularities*, if $R^i f_*(\mathcal{O}_Y) = 0$ for some desingularization $f : Y \rightarrow X$ and for any $i \geq 1$; then X is Cohen-Macaulay with dualizing sheaf $\mathcal{O}(K_X)$.

Proposition 2.4. *Let X be an affine G -variety, with degeneration X_0 corresponding to the choice of a reduced decomposition of w_0 . Then X is normal (resp. has rational singularities) if and only if X_0 is normal (resp. has rational singularities). Further, if $mK_X \sim 0$ for some integer m , then $mK_{X_0} \sim 0$.*

Proof. If X_0 is normal, then so is X e.g. by [Gro97, Thm.18.3]. Conversely, if $X = \operatorname{Spec}(R)$ is normal, then so is $\operatorname{Spec}(R^U)$ by [Gro97, Thm.18.4]. It follows easily that $X_0 = \operatorname{Spec}(R^U \otimes A)^T$ is normal as well.

The assertion on rational singularities is proved similarly, by using the stability of rational singularities under deformations [Elk78] and under quotients by reductive groups [Bou87].

Now assume that X (or X_0) is normal ; then $\mathcal{X}$ is normal by Serre's criterion. If $mK_X \sim 0$, then the sheaf $\mathcal{O}(mK_X)$ is isomorphic to $\mathcal{O}_X$ outside X_0 . Since both sheaves are divisorial, and X_0 is irreducible, it follows that $\mathcal{O}(mK_X) \cong \mathcal{O}_X(nX_0)$ for some integer n . But X_0 is the divisor of the regular function π on $\mathcal{X}$, so that $\mathcal{O}(mK_X) \cong \mathcal{O}_X$. Restricting to X_0 , we obtain $\mathcal{O}(mK_{X_0}) \cong \mathcal{O}_{X_0}$. Indeed, it suffices to check this on the regular locus of X_0 ; the latter is contained in the regular locus of $\mathcal{X}$, where the adjunction formula $\mathcal{O}(mK_X)|_{X_0} \cong \mathcal{O}(mK_{X_0})$ holds. $\square$

Remarks 2.5. (i) Since toric varieties have rational singularities, Propositions 2.3 and 2.4 yield another proof of the rationality of singularities of spherical varieties ([Pop87, Thm.10]).

(ii) For any reduced decomposition of w_0 , the limit $(G//U)_0$ has trivial canonical class. Indeed, $G//U$ contains the homogeneous space G/U as an open orbit, with complement of codimension at least 2. Further, $K_{G/U} \sim 0$, since U acts by unimodular transformations on the tangent space to G/U at the base point. Thus, Proposition 2.4 applies.

On the other hand, $(G//U)_0$ is singular if (say) the group G is simple of rank ≥ 2 . Indeed, $G//U$ is singular in that case.

3. DEGENERATIONS OF POLARIZED PROJECTIVE G -VARIETIES

Definition 3.1. A *polarized (projective) G -variety* is a pair (X, L) , where X is a normal projective variety equipped with a G -action, and L is an ample G -linearized invertible sheaf on X .

For any integer n , the sheaf $L^n := L^{\otimes n}$ is also G -linearized. Thus, the space $H^0(X, L^n)$ is a finite-dimensional rational G -module. Further, the graded algebra

$$R(X, L) := \bigoplus_{n=0}^{\infty} H^0(X, L^n)$$

is a finitely generated, integrally closed domain. We have: $X = \text{Proj } R(X, L)$, and $L^n = \mathcal{O}_X(n)$ for all integers n .

Next we choose a reduced decomposition $\underline{w}_0$ of w_0 , which yields the associated graded algebra, $\text{gr } R(X, L)$. By Proposition 2.4, this algebra is still a finitely generated, integrally closed domain, with an action of $\mathbb{C}^* \times T \times \mathbb{T}$ such that the $\mathbb{C}^*$ -action defines a positive grading. Thus,

$$X_0 := \text{Proj gr } R(X, L)$$

is a projective $T \times \mathbb{T}$ -variety equipped with $T \times \mathbb{T}$ -linearized sheaves

$$L_0^{(n)} := \mathcal{O}_{X_0}(n)$$

for all integers n . By [Dem88, Thm.3.5], there exists a $\mathbb{Q}$ -Weil divisor D on X_0 such that $L_0^{(n)} \cong \mathcal{O}_{X_0}(nD)$ for all n . (Here $\mathcal{O}_{X_0}(nD)$ denotes the sheaf of rational functions f on X_0 such that the $\mathbb{Q}$ -Weil divisor $(f) + nD$ has non-negative coefficients). Further, D is $\mathbb{Q}$ -Cartier and ample, i.e., the sheaf $L_0^{(m)}$ is invertible and ample for any sufficiently divisible integer $m > 0$. In particular, every sheaf $L_0^{(n)}$ is divisorial, i.e., it is the sheaf of sections of an integral Weil divisor (called the floor of nD).

We then say that (X, L) *degenerates to the $\mathbb{Q}$ -polarized variety (X_0, L_0)* , and that (X_0, L_0) *is a limit of (X, L)* . (To be completely correct, we should say that

(X, L^n) degenerates to $(X_0, L_0^{(n)})$ for all positive integers n , as L_0 does not determine uniquely all the $L_0^{(n)}$. Since $\text{gr } R(X, L) \cong R(X, L)$ as a $\mathbb{C}^* \times T$ -module, each space $H^0(X_0, L_0^{(n)})$ is isomorphic to $H^0(X, L^n)$ as a T -module, but it acquires a compatible action of the torus $\mathbb{T}$.

Now Propositions 2.2 and 2.3 may be generalized as follows.

Theorem 3.2. *Let (X, L) be a polarized G -variety and choose a reduced decomposition $\underline{w}_0$. Then there exists a family of T -varieties $\pi : \mathcal{X} \rightarrow \mathbb{A}^1$, where $\mathcal{X}$ is a normal variety, together with divisorial sheaves $\mathcal{O}_{\mathcal{X}}(n)$ ($n \in \mathbb{Z}$), such that*

- (i) π is projective and flat.
- (ii) π is trivial with fiber X over the complement of 0 in $\mathbb{A}^1$, and $\mathcal{O}_{\mathcal{X}}(n)|_X \cong L^n$ for all n .
- (iii) The fiber of π at 0 is isomorphic to X_0 , and $\mathcal{O}_{\mathcal{X}}(n)|_{X_0} \cong L_0^{(n)}$ for all n .

If, in addition, X is spherical, then X_0 is a toric variety under $T \times \mathbb{T}$.

Proof. Let $R := R(X, L)$ with filtration $(R_{\leq m})$ and let $\mathcal{R} := \bigoplus_{m=0}^{\infty} R_{\leq m} t^m$, as in the proof of Proposition 2.2. Then the Rees algebra $\mathcal{R}$ carries two compatible $\mathbb{N}$ -gradings: by m , and by the degree inherited from that of R . For the second grading, the subspace $\mathcal{R}_0$ is just the polynomial ring $\mathbb{C}[t]$. Put $\mathcal{X} := \text{Proj } \mathcal{R}$ with corresponding twisting sheaves $\mathcal{O}_{\mathcal{X}}(n)$. Then $\mathcal{X}$ is equipped with a morphism to $\mathbb{A}^1$ such that the assertions (i), (ii) and (iii) are easily checked.

Since R and $\text{gr } R(X, L)$ are normal, it follows that $\mathcal{R}$ is normal, by Serre's criterion. So $\mathcal{X}$ is normal as well. Consider the variety $\hat{\mathcal{X}} := \text{Spec}(\mathcal{R})$ and the morphism $t : \hat{\mathcal{X}} \rightarrow \mathbb{A}^1$. The latter admits a section, corresponding to the projection $\mathcal{R} \rightarrow \mathcal{R}_0 = \mathbb{C}[t]$. We still denote by $\mathbb{A}^1 \subset \hat{\mathcal{X}}$ the image of this section. Now we have a morphism

$$p : \hat{\mathcal{X}} - \mathbb{A}^1 \rightarrow \mathcal{X}$$

which is a geometric quotient by the $\mathbb{C}^*$ -action corresponding to the second grading. Further, using the normality of $\hat{\mathcal{X}}$, one checks that each $\mathcal{O}_{\mathcal{X}}(n)$ is the eigenspace of degree n in the sheaf $p_* \mathcal{O}_{\hat{\mathcal{X}} - \mathbb{A}^1}$. This sheaf is reflexive, as $\hat{\mathcal{X}} - \mathbb{A}^1$ is normal and p is equidimensional. Thus, all the $\mathcal{O}_{\mathcal{X}}(n)$ are reflexive as well. Since they have generic rank 1, it follows that they are divisorial. $\square$

Next recall that polarized toric varieties correspond to integral convex polytopes, and that Fano toric varieties correspond to reflexive polytopes, see e.g. [Bat94]. The assignment of a polytope to a polarized toric variety may be generalized to arbitrary polarized projective G -varieties by the following well-known result.

Lemma 3.3. *Let (X, L) be a polarized G -variety. Then the points $\frac{\lambda}{n} \in \Lambda_{\mathbb{Q}}$ such that: $\lambda \in \Lambda^+$, n is a positive integer, and the isotypical component $H^0(X, L^n)_{(\lambda)}$ is nonzero, are exactly the rational points of a rational convex polytope $P(X, L)$ of $\Lambda_{\mathbb{R}}$. Further, $P(X, L^m) = mP(X, L)$ for any positive integer m .*

Indeed, $H^0(X, L^n)_{(\lambda)} \neq 0$ if and only if $H^0(X, L^{(n)})_{\lambda}^U \neq 0$. Further, the algebra $R(X, L)^U$ is finitely generated and $\Lambda^+ \times \mathbb{N}$ -graded; let (f_i) be homogeneous generators, and (λ_i, n_i) their degrees. Then one easily checks that $P(X, L)$ is the rational convex hull of the points $\frac{\lambda_i}{n_i}$.

Definition 3.4. $P(X, L)$ is the *moment polytope* of the polarized G -variety (X, L) .

By positive homogeneity of the moment polytope, this definition extends to $\mathbb{Q}$ -polarized varieties, in particular, to any limit (X_0, L_0) of (X, L) . We denote by $Q_{\underline{w}_0}(X, L)$ the moment polytope of that limit. It is a rational convex polytope in $\Lambda_{\mathbb{R}} \times \mathbb{R}^N$, related to the moment polytope of (X, L) by the following

Theorem 3.5. *The first projection $p : \Lambda_{\mathbb{R}} \times \mathbb{R}^N \rightarrow \Lambda_{\mathbb{R}}$ restricts to a surjective map*

$$p : Q_{\underline{w}_0}(X, L) \rightarrow P(X, L),$$

with fiber over any $\lambda \in \Lambda_{\mathbb{R}}^+$ being the string polytope $Q_{\underline{w}_0}(\lambda^*)$. In particular, for $\lambda \in \Lambda^+$, the limit of the flag variety G/P_{λ} is a toric variety under $\mathbb{T}$, and its moment polytope is the string polytope $Q_{\underline{w}_0}(\lambda^*)$.

Proof. By definition, the rational points of $Q_{\underline{w}_0}(X, L)$ are the pairs $(\frac{\lambda}{n}, \frac{\varphi}{n})$ such that: $H^0(X, L^n)_{(\lambda)} \neq 0$, and $(\lambda^*, \varphi) \in \mathcal{C}_{\underline{w}_0}$. This implies the first assertion for rational λ , and hence for all λ since all involved polytopes are rational.

Consider the G -variety $X = G/P_{\lambda}$ and the G -linearized line bundle $L = L_{\lambda}$. Then L is ample, and $R(X, L) = \bigoplus_{n=0}^{\infty} V(n\lambda^*)$. Thus, X is spherical with moment polytope being the point λ^* . This implies the second assertion. $\square$

Remark 3.6. The moment polytope $P(X, L)$ need not be integral. Consider, for example, the group GL_n acting on the projectivization of the space of $n \times n$ matrices by left multiplication. Then one checks that the moment polytope for the line bundle $\mathcal{O}(1)$ has nonintegral vertices as soon as $n \geq 3$.

Finally, we consider limits of polarized spherical varieties which are $\mathbb{Q}$ -Fano in the sense of the following

Definition 3.7. We say that a normal projective variety X is a $\mathbb{Q}$ -Gorenstein Fano (or simply $\mathbb{Q}$ -Fano) if its anticanonical class $-K = -K_X$ is $\mathbb{Q}$ -Cartier and ample. If $-K_X$ is Cartier and ample, we say that X is (Gorenstein) Fano.

Theorem 3.8. *Let (X, L) be a polarized spherical G -variety with limit (X_0, L_0) . If X is a $\mathbb{Q}$ -Fano and $L = \mathcal{O}(-mK_X)$ for some positive integer m , then X_0 is a $\mathbb{Q}$ -Fano and $L_0^{(n)} = \mathcal{O}(-nmK_{X_0})$ for all n .*

Further, $L_0^{(n)}$ is invertible if and only if the polytope $nQ(X, L)$ is integral; in this case, $nP(X, L)$ is integral as well.

In particular, if X is a Fano and $L = \mathcal{O}(-K_X)$, then $L_0 = \mathcal{O}(-K_{X_0})$. Thus, X_0 is a Fano if and only if $Q(X, L)$ is integral; in this case, $Q(X, L)$ is reflexive.

Proof. The isomorphism $L_0^{(n)} = \mathcal{O}(-nmK_{X_0})$ is proved by the argument of Proposition 2.4. It implies, of course, that X_0 is $\mathbb{Q}$ -Fano.

If $nQ(X, L)$ is integral, then its vertices yield global sections of $\mathcal{O}_{X_0}(n)$, eigenvectors of the torus $\mathbb{T}$. These sections have no common zeroes (since they do not vanish simultaneously at any $\mathbb{T}$ -fixed point), so that $\mathcal{O}_{X_0}(n)$ is invertible. Conversely, if $\mathcal{O}_{X_0}(n)$ is invertible, then $-nmK$ is an ample (integral) Cartier divisor on the toric variety X_0 . Thus, it is generated by its global sections, and the preceding argument shows that all the vertices of $nQ(X, L)$ are integral. This proves the second assertion. $\square$

Remarks 3.9. (i) For an arbitrary polarized G -variety (X, L) with limit (X_0, L_0) , the following conditions are equivalent:

(a) X has rational singularities and $H^i(X, L^n) = 0$ for all $i \geq 1$ and $n \geq 0$.

(b) X_0 has rational singularities and $H^i(X_0, L_0^{(n)}) = 0$ for all $i \geq 1$ and $n \geq 0$.

Indeed, one checks that (a) (resp. (b)) is equivalent to the rationality of singularities of $\text{Spec } R(X, L)$ (resp. $\text{Spec } R(X_0, L_0)$), and one applies Proposition 2.4.

(ii) By the Kawamata-Viehweg vanishing theorem, the preceding assumption (a) holds in the case where X is a $\mathbb{Q}$ -Fano with log terminal singularities, and $L = \mathcal{O}(-mK_X)$. Together with the argument of Proposition 2.4, it follows that the limit X_0 is a $\mathbb{Q}$ -Fano, and $L_0^{(n)} = \mathcal{O}(-mnK_{X_0})$ for all n . This applies, for example, to Fano varieties with their anticanonical polarization.

(iii) For a polarized spherical variety (X, L) , one has $H^i(X, L^n) = 0$ for all $i \geq 1$ and $n \geq 0$ (as follows e.g. from the rationality of singularities of $\text{Spec } R(X, L)$). Thus, the Hilbert function $n \mapsto h^0(X, L^n)$ is a polynomial for $n \geq 0$, and coincides with the Ehrhart function

$$n \mapsto \#(Q_{\underline{w}_0}(X, L^n) \cap \mathbb{Z}^N) = \#(nQ_{\underline{w}_0}(X, L) \cap \mathbb{Z}^N)$$

of the polytope $Q_{\underline{w}_0}(X, L)$. This generalizes a result of Okounkov [Oko97].

(iv) Let X be a Fano spherical variety and $L = \mathcal{O}(-K_X)$. If the limit X_0 is nonsingular, then it is isomorphic to X as a T -variety. Indeed, X_0 is Fano by Theorem 3.8. Since X_0 is toric, it follows easily that $H^1(X_0, T_{X_0}) = 0$. This implies, in turn, our rigidity statement by the argument of [AB04, Thm.3.1].

As a consequence, any anticanonical limit of a flag variety $X = G/P$ is singular, except if X is a product of projective spaces.

4. STRING AND MOMENT POLYTOPES

Recall that $p : \Lambda_{\mathbb{R}} \times \mathbb{R}^N \rightarrow \Lambda_{\mathbb{R}}$ denotes the projection onto the first summand; it maps the cone $\mathcal{C}_{\underline{w}_0}$ onto the positive Weyl chamber $\Lambda_{\mathbb{R}}^+$.

Definition 4.1. For every $\lambda \in \Lambda_{\mathbb{R}}$, let σ_{λ}^0 be the intersection of the images $p(\tau^0)$, where τ is a face of the cone $\mathcal{C}_{\underline{w}_0}$ with relative interior τ^0 , and $p(\tau^0)$ contains λ . Let $\Sigma_{\underline{w}_0}$ be the collection of the closed cones $\sigma_{\lambda} := \overline{\sigma_{\lambda}^0}$.

Since the intersection of any two faces of $\mathcal{C}_{\underline{w}_0}$ is a face, the intersection of any two cones of $\Sigma_{\underline{w}_0}$ is again in $\Sigma_{\underline{w}_0}$. Hence, $\Sigma_{\underline{w}_0}$ is a fan, with support $\Lambda_{\mathbb{R}}^+$. The following Lemma follows by an elementary convexity argument.

Lemma 4.2. *Two weights $\lambda, \mu \in \Lambda_{\mathbb{R}}^+$ lie in the same cone of $\Sigma_{\underline{w}_0}$ if and only if the polytope $Q_{\underline{w}_0}(\lambda + \mu)$ is the Minkowski sum of the polytopes $Q_{\underline{w}_0}(\lambda)$ and $Q_{\underline{w}_0}(\mu)$.*

Corollary 4.3. *If the fan $\Sigma_{\underline{w}_0}$ is trivial, i.e. consists of the positive chamber $\Lambda_{\mathbb{R}}^+$ and its faces, then for all regular dominant weights λ the polytopes $Q_{\underline{w}_0}(\lambda)$ have the same normal fan. In this case, the polytope $Q_{\underline{w}_0}(X, L)$ is integral if and only if the string polytope $Q_{\underline{w}_0}(\lambda)$ is integral for any vertex λ of $P(X, L)$.*

If the fan $\Sigma_{\underline{w}_0}$ is non-trivial then clearly the vertices of $Q_{\underline{w}_0}(X, L)$ are among the vertices of the polytopes $Q_{\underline{w}_0}(\lambda)$ as λ goes over the vertices in the subdivision $\Sigma_{\underline{w}_0} \cap P(X, L)$ of $P(X, L)$.

By Theorem 3.8, the anticanonical limit of the flag variety G/B is a Fano if and only if the polytope $Q_{\underline{w}_0}(G/B, \mathcal{O}(-K)) = Q_{\underline{w}_0}(2\rho)$ is integral, where ρ is the half-sum of positive roots. This is not true in general, as we show below in Example 5.11. However, we will show that any toric limit (X_0, L_0) of $(G/P_{\lambda}, L_{\lambda})$ is Gorenstein at its $\mathbb{T}$ -fixed points associated with extremal weight vertices (in the

sense of Definition 1.3). Recall that any such vertex q_μ is integral, and hence defines a global section s_μ of L_0 , eigenvector of $\mathbb{T}$ of weight v_μ . The complement of the zero locus of s_μ is a $\mathbb{T}$ -stable affine open subset of X_0 , denoted by $X_{0,\mu}$.

Theorem 4.4. *For every extremal weight vertex q_μ of $Q_{\underline{w}_0}(\lambda)$, the corresponding open subset $X_{0,\mu}$ has trivial canonical class.*

Proof. Let $\pi : \mathcal{X} \rightarrow \mathbb{A}^1$ and $\mathcal{O}_{\mathcal{X}}(1)$ be as in Proposition 3.2. The extremal weight vertex $q_\mu = \pi_\lambda^{-1}(\mu)$ defines a 1-dimensional submodule of $\pi_*\mathcal{O}_{\mathcal{X}}(1)$ and hence a global section $s_\mu \in H^0(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(1))$. The family $\mathcal{X}_\mu = \{s_\mu \neq 0\} \rightarrow \mathbb{A}^1$ is an affine family whose fiber at any nonzero point is the complement of the zero locus in G/P_λ of the extremal weight vector $v_\mu \in H^0(G/P_\lambda, L_\lambda)$. This fiber is a W -translate of the negative cell U^-P_λ/P_λ ; in particular, it is an affine space. Hence, it has trivial canonical class, so that the same holds for $X_{0,\mu}$ by the proof of Proposition 2.4. $\square$

Next recall that a dominant weight λ is called *minuscule* if the only weights of $V(\lambda)$ are the extremal weights $\mu \in W\lambda$; equivalently, the only integral points in $Q_{\underline{w}_0}(\lambda)$ are the extremal weight vertices. Clearly, the set of minuscule weights is stable under $\lambda \mapsto \lambda^*$. All the minuscule weights are fundamental; their list is as follows:

- Type A_n : any fundamental weight.
- Type B_n : ω_n .
- Type C_n : ω_1 .
- Type D_n : $\omega_1, \omega_{n-1}, \omega_n$.
- Type E_6 : ω_1, ω_6 .
- Type E_7 : ω_7 .
- Types G_2, F_4, E_8 : none.

Also recall that a fundamental weight λ is called *cominuscule* if the corresponding fundamental weight in the dual root system is minuscule. The cominuscule weights are, of course, the minuscule weights in type A, D and E , plus the following:

- Type B_n : ω_1 .
- Type C_n : ω_n .

Thus, the flag varieties G/P_λ associated with the (co)minuscule weights are the grassmanians, the quadrics, the grassmanians of maximal isotropic subspaces (with respect to a symmetric or alternating non degenerate bilinear form), and two exceptional varieties.

Theorem 4.5. *For any (co)minuscule weight λ and for any reduced decomposition $\underline{w}_0$, the polytope $Q_{\underline{w}_0}(\lambda)$ is integral. Further, the corresponding toric variety $(G/P_\lambda)_0$ is a Fano. In the minuscule case, all the vertices of $Q_{\underline{w}_0}(\lambda)$ are extremal weight vertices.*

Proof. We begin with the minuscule case. For any positive integer n , we show that the dual canonical basis of $V(n\lambda^*)$ coincides (up to nonzero scalar multiples) with its *standard monomial basis*. The latter can be defined as follows (see e.g. [GL96]). Let W^λ be the set of representatives of minimal length of all cosets wW_λ , where W_λ is the isotropy group of λ in W . Then we may index the extremal weight vectors of $V(\lambda^*)$ by W^λ , via

$$x \mapsto p_x := v_{-x(\lambda)} \in H^0(G/P_\lambda, L_\lambda) = V(\lambda^*).$$

The set W^λ is partially ordered by the Bruhat ordering of W . Now the products

$$p_{x_1} \cdots p_{x_n} \in H^0(G/P_\lambda, L_\lambda^n) = V(n\lambda^*),$$

where $x_1 \geq \cdots \geq x_n$ in W^λ , form a basis of $V(n\lambda^*)$. These products are called the standard monomials. Further, if $x \geq y \in W^\lambda$, then there exists a sequence $(s_{i_1}, \dots, s_{i_\ell})$ of simple reflections such that

$$x = s_{i_1} \cdots s_{i_\ell} y \quad \text{and} \quad \ell(x) = \ell + \ell(y),$$

as follows from [GL96, Lem.7.1]; we say that the Bruhat ordering of W^λ is generated by simple reflections. Thus, given $x_1 \geq \cdots \geq x_n$ in W^λ , there exists a reduced decomposition $w_0 = s_{i_1} \cdots s_{i_N}$ such that each x_k is a subword $s_{i_1} \cdots s_{i_j}$ for some $j = j(k)$. Now we may apply [CL02, Thm.14] which asserts that a nonzero scalar multiple of $p_{x_1} \cdots p_{x_n}$ lies in the dual canonical basis of $V(n\lambda^*)$.

Together with the multiplicative property of this basis, it follows that any integral point of $Q_{\underline{w}_0}(n\lambda)$ is a sum of n extremal weight vertices. Since $Q_{\underline{w}_0}(n\lambda) = nQ_{\underline{w}_0}(\lambda)$, it follows in turn that the polytope $Q_{\underline{w}_0}(\lambda)$ is integral and that its integral points are precisely the extremal weight vertices.

Now, if λ is cominuscule, then it is of classical type in the sense of [LS86]. Then a basis for $V(\lambda^*)$ consists of T -eigenvectors $p_{x,y}$, where $x, y \in W^\lambda$ form an admissible pair in the sense of [loc.cit.]; in particular, $x \geq y$ (and $x = y$ yields the extremal weight vectors). Further, by [loc.cit.] again, a basis of $V(n\lambda^*)$ consists of the standard monomials of degree n in the $p_{x,y}$, i.e., of those products $p_{x_1,y_1} \cdots p_{x_n,y_n}$ such that $x_1 \geq y_1 \geq \cdots \geq x_n \geq y_n$. Since the Bruhat ordering in W^λ is still generated by simple reflections, [CL02, Thm.14] also applies and yields that the dual canonical basis of $V(n\lambda^*)$ coincides (up to nonzero scalar multiples) with its standard monomial basis. As above, it follows that any integral point of $Q_{\underline{w}_0}(n\lambda)$ is a sum of n integral points of $Q_{\underline{w}_0}(\lambda)$. This implies, in turn, that the polytope $Q_{\underline{w}_0}(\lambda)$ is integral.

It remains to show that $(G/P_\lambda)_0$ is a Fano. Note that $\mathcal{O}(-K_{G/P_\lambda}) = L_{m\lambda}$ for some positive integer m (as the weight λ is fundamental), so that the polytope $Q_{\underline{w}_0}(G/P_\lambda, \mathcal{O}(-K)) = mQ_{\underline{w}_0}(\lambda)$ is integral. Now Theorem 3.8 completes the proof. $\square$

5. EXAMPLES AND COUNTEREXAMPLES

§ 5.1. G of type A_n and $\underline{w}_0^{\text{std}}$ (Gelfand-Tsetlin case). The simplest reduced decomposition of the longest element in $W = S_{n+1}$ is

$$\underline{w}_0^{\text{std}} = (s_1)(s_2s_1)(s_3s_2s_1) \cdots (s_ns_{n-1} \cdots s_1),$$

where s_i denotes the transposition exchanging i with $i+1$. It is convenient to use coordinates $x_{i,j}$ with $i, j \geq 1$, $i+j \leq n+1$ in place of $t_1, \dots, t_{n(n+1)/2}$. The string cone is defined by

$$x_{n,1} \geq 0; \quad x_{n-1,2} \geq x_{n-1,1} \geq 0; \quad \dots \quad x_{1,n} \geq \cdots \geq x_{1,1} \geq 0,$$

and the remaining λ -inequalities are

$$x_{i,j} \leq \langle \lambda, \alpha_j^\vee \rangle - x_{i,j-1} + \sum_{k=1}^{i-1} (-x_{k,j-1} + 2x_{k,j} - x_{k,j+1})$$

for all $k = 1, \dots, n$, $i = 1, \dots, k$ in which we set a variable x to zero whenever its indices are out of bounds.


The corresponding polytopes $Q_{\underline{w}_0^{\text{std}}}(\lambda)$ are known as *Gelfand-Tsetlin polytopes*, and we will use the notation $\text{GT}(\lambda)$ for them. A more familiar description of $\text{GT}(\lambda)$ is in terms of *Gelfand-Tsetlin patterns*

$$g_{i,j} \geq g_{i+1,j} \geq g_{i,j+1}$$

in variables $g_{i,j}$, $i, j \geq 1$, $i + j \leq n + 1$ and with variables $g_{0,i}$ corresponding to λ :

$$\lambda = \sum_{i=0}^{n+1} g_{0,i} \epsilon_i = \sum_{i=0}^{n+1} \lambda_i \epsilon_i, \quad \text{so that} \quad \langle \lambda, \alpha_j^\vee \rangle = \langle \lambda, \epsilon_j - \epsilon_{j+1} \rangle = \lambda_j - \lambda_{j+1}.$$

The inequalities can be represented by a graph Γ in which an arrow goes from a smaller variable to a larger:



The linear change of coordinates is given by

$$x_{i,j} = \sum_{k=1}^j (g_{i-1,k} - g_{i,k}), \quad g_{i,j} = \lambda_j + \sum_{k=1}^i (x_{k,j-1} - x_{k,j}).$$

Lemma 5.1. *The fan $\Sigma_{\underline{w}_0^{\text{std}}}$ is trivial.*

Proof. Indeed, the additive property $\text{GT}(\lambda + \mu) = \text{GT}(\lambda) + \text{GT}(\mu)$ is obvious from the Gelfand-Tsetlin patterns. (This was already noticed by Kaveh in [Kav03].) $\square$

Lemma 5.2. $\text{GT}(\lambda)$ is integral $\iff \langle \lambda, \alpha_i^\vee \rangle \in \mathbb{Z}$ for all i .

Proof. The vertices are given by setting some of the Gelfand-Tsetlin inequalities to equalities. It is then clear that $g_{i,j}$ are going to be equal to some λ_j 's and that all of them appear. So if the λ_j 's are integral then so are the $g_{i,j}$'s and vice versa. $\square$

In fact, the following explicit description of the vertices is obvious:

Lemma 5.3. *Each vertex of the Gelfand-Tsetlin polytope $\text{GT}(\lambda)$ corresponds to setting each variable $g_{i,j}$ to some λ_j in the way allowed by inequalities. Equivalently, the boundaries between different values of $g_{i,j}$'s are paths Π_i from the boundary of the graph Γ along the diagonal to the lower left corner such that the union $\cup \Pi_i$ is a tree.*

The collections $\cup \Pi_i$ were called *meanders* in [BCFKvS00] in which the authors describe the toric degeneration constructed by Gonciulea and Lakshmibai. Note that in the case when the weight λ is not regular, i.e. when some $\lambda_j = 0$ and the corresponding variety G/P_λ is a partial flag variety, the graph Γ can be simplified by cutting out square corners along the diagonal in which the values of $g_{i,j}$'s are uniquely determined.

The extremal weight vertices of $\text{GT}(\lambda)$ can be described with a little more work. Since we do not need this description, we leave the proof to the reader.

Lemma 5.4. *For every permutation $w \in W = S_{n+1}$, $w = (k_1 k_2 \dots k_{n+1})$ the corresponding vertex is obtained by filling $k_1 - 1$ variables $g_{i,j}$ starting from the top in a minimal way with λ_1 , then filling $k_2 - 1$ variables in a minimal way with λ_2 etc.*

Example 5.5. For the group $G = \text{SL}_3$ of type A_2 and a regular weight λ , there are $6 = 3!$ extremal weight vertices and one additional vertex corresponding to setting $g_{1,1} = g_{1,2} = g_{2,1} = \lambda_2$. The polytope $\text{GT}(\lambda)$ has its projection to $\Lambda_{\mathbb{R}}$ shown on Fig. 1.




FIGURE 1. Projection of $\text{GT}(\lambda)$

§ 5.2. G of type A_n and other $\underline{w}_0$. For many of the explicit computations below, we used the freely available program `porta` for performing manipulations with polytopes.

The fans $\Sigma_{\underline{w}_0}$ need not be trivial in general.

Example 5.6. Let G be of type A_3 and let the reduced decomposition be $\underline{w}_0 = s_1 s_3 s_2 s_3 s_1 s_2$. The fan $\Sigma_{\underline{w}_0}$ consists of two maximal dimensional cones obtained by splitting $\Lambda_{\mathbb{R}}^+$ into two halves by the hyperplane $\langle \lambda, \alpha_1^\vee \rangle = \langle \lambda, \alpha_3^\vee \rangle$. For a regular weight λ , the polytope $Q_{\underline{w}_0}(\lambda)$ has 38 or 44 vertices depending on which cone λ lies in.

Example 5.7. The polytopes $Q_{\underline{w}_0}(\rho)$ are integral for $G = \text{SL}_n$, $n \leq 5$. In type A_3 they have 12 or 13 facets and 38, 40 or 42 vertices. In type A_4 , the polytopes have from 20 to 27 facets and from 334 to 425 vertices.

Conjecture 5.8. *For G of type A_n and any reduced decomposition $\underline{w}_0$, the polytope $Q_{\underline{w}_0}(\lambda)$ is integral if and only if $\langle \lambda, \alpha_i^\vee \rangle \in \mathbb{Z}$ for all i .*

§ 5.3. Polytopes for fundamental weights.

Example 5.9. A direct computation shows that for a group of type A_n , $n \leq 4$, the polytopes $Q_{\underline{w}_0}(\omega_i)$ for a fixed fundamental weight ω_i are isomorphic, i.e. they do not depend on $\underline{w}_0$. However, they are different as polytopes marked by two distinguished vertices (the highest and lowest weight vertices). For example, in type A_3 the tangent cone at the origin of the polytope $Q_{\underline{w}_0^{\text{std}}}(\omega_2)$ is non-simplicial. On the other hand, for $\underline{w}_0 = s_2 s_1 s_2 s_3 s_2 s_1$ the tangent cone of $Q_{\underline{w}_0}(\omega_2)$ at the origin is simplicial.

§ 5.4. Other classical types.

Example 5.10. For G of type C_2 , Littelmann [Lit98, §2] computes the string cones for the reduced decompositions $\underline{w}_0 = s_1 s_2 s_1 s_2$ and $\underline{w}'_0 = s_2 s_1 s_2 s_1$. They are respectively

$$\{t_1 \geq 0; \quad 2t_2 \geq t_3 \geq 2t_4 \geq 0\} \quad \text{and} \quad \{t_1 \geq 0; \quad t_2 \geq t_3 \geq t_4 \geq 0\}.$$

In both cases, the fan Σ is trivial. A simple computation shows that in the second case all the polytopes $Q(\lambda)$ are integral. However, in the first case, the polytope $Q(\lambda)$ is integral if and only if $\langle \lambda, \alpha_1^\vee \rangle$ is even. In particular, $Q(2\rho) = Q(G/B, \mathcal{O}(-K))$ is integral. The polytopes $Q_{\underline{w}_0}(2\rho)$ and $Q_{\underline{w}'_0}(2\rho)$ both have 12 vertices.

There is a notion of Gelfand-Tsetlin polytopes in types B , C and D . Their defining inequalities can be found in [BZ88] and [Lit98, §6, §7]. From the definitions, it is obvious that in types B_n and C_n the fan $\Sigma_{\underline{w}_0^{\text{std}}}$ is trivial and the polytope $Q_{\underline{w}_0^{\text{std}}}(\lambda)$ is integral whenever the $\langle \lambda, \alpha_i^\vee \rangle$ are all even. In particular, the polytope $Q_{\underline{w}_0^{\text{std}}}(2\rho)$ for the anticanonical limit of G/B is (integral and) reflexive, and the corresponding variety $(G/B)_0$ is a Fano.

In type D_n , the fan $\Sigma_{\underline{w}_0^{\text{std}}}$ is non-trivial.

§ 5.5. G of exceptional type.

Example 5.11. For G of type E_6 and some nice decomposition $\underline{w}_0$, the polytope $Q_{\underline{w}_0}(2\rho) = Q_{\underline{w}_0}(G/B, \mathcal{O}(-K))$ is not integral.

Littelmann [Lit98, §8] computed (and we checked) the string cone for the reduced decomposition $\underline{w}_0 = \tau \cdot s_6 s_2 s_3 s_1 s_4 s_5 s_3 s_4 s_2 s_3 s_1 s_6 s_2 s_3 s_4 s_5$ with the enumeration

$$\begin{array}{ccccccc} & & & \alpha_1 & & & \\ & & & | & & & \\ \alpha_5 & \text{---} & \alpha_4 & \text{---} & \alpha_3 & \text{---} & \alpha_2 & \text{---} & \alpha_6 \end{array}$$

of the simple roots, where τ is a reduced decomposition for the longest element in the Weyl group of the sub-root system of type D_5 obtained by suppressing α_6 . The cone $C_{\underline{w}_0}$ is the product of the string cone for D_5 and the cone

$$\begin{aligned} t_1 \geq t_2 \geq t_3 \geq \begin{Bmatrix} t_4 \\ t_5 \end{Bmatrix} \geq t_7 \geq \begin{Bmatrix} t_8 \\ t_9 \end{Bmatrix} \geq t_{10} \geq \begin{Bmatrix} t_{11} \\ t_{13} \end{Bmatrix} \geq t_{14} \geq t_{15} \geq t_{16} \geq 0; \\ t_5 \geq t_6 \geq t_8 \geq 0; \quad t_9 \geq t_{12} \geq t_{13} \geq 0. \end{aligned}$$

Hence, the projection of the 36-dimensional polytope $Q_{\underline{w}_0}(\lambda)$ to the last 16 coordinates is the 16-dimensional polytope Q' defined by the inequalities above and the λ -inequalities for $k = 36, 35, \dots, 21$. An explicit computation shows that for $\lambda = \rho$,

i.e. when all $\langle \lambda, \alpha_j^\vee \rangle = 1$, the polytope nQ' is integral if and only if n is divisible by 6. Hence, the polytopes $2Q'$ and $Q_{\underline{w}_0}(2\rho)$ are not integral, and the limit toric variety of the anticanonically polarized variety G/B is not a Fano.

§ 5.6. String polytopes and Duistermaat-Heckman measure. An important question is what the string polytopes $Q_{\underline{w}_0}(\lambda)$ have in common for different reduced decompositions $\underline{w}_0$. Here, we note two such properties: the Ehrhart function

$$n \mapsto \#(Q_{\underline{w}_0}(n\lambda) \cap \mathbb{Z}^N) = \dim V(n\lambda^*) = \dim V(n\lambda)$$

for $n \geq 0$, which is a polynomial function even if $Q_{\underline{w}_0}(\lambda)$ is not integral, and the Duistermaat-Heckman measure.

Recall that for any symplectic (differentiable) manifold (X, ω) of dimension $2n$, equipped with a Hamiltonian action of a compact torus K , the Duistermaat-Heckman measure $d\sigma_{DH}^K$ is the pushforward of the Liouville measure $\omega^n/n!$ under the moment map $\mu : X \rightarrow (\text{Lie } K)^*$ to the dual of the Lie algebra of K . The image $\mu(X)$ is a convex polytope, and the measure $d\sigma_{DH}^K$ is continuous and piecewise polynomial on this polytope.

On the other hand, let (X, L) be a (possibly singular) polarized projective toric variety under the torus $\mathbb{T}$. Denote by $\mathbb{K} \subset \mathbb{T}$ the maximal compact subtorus. Then X admits a moment map $\mu_{\mathbb{K}} : X \rightarrow (\text{Lie } \mathbb{K})^*$, which may be defined as the composition of the finite morphism $X \rightarrow \mathbb{P}H^0(X, L)^*$, with the moment map for the $\mathbb{K}$ -action on $\mathbb{P}H^0(X, L)^*$ (regarded as a symplectic variety via a $\mathbb{K}$ -invariant Kähler structure). The image of the moment map identifies with the moment polytope $P(X, L)$, and the measure $d\sigma_{DH}^{\mathbb{K}}$ is just the standard Euclidean measure multiplied by $(2\pi)^{\dim X}$ on this polytope.

Given another torus T with maximal compact subtorus K , and a homomorphism $h : T \rightarrow \mathbb{T}$, the K -action on X via h admits a moment map $\mu_K : X \rightarrow (\text{Lie } K)^*$ which is nothing but the map $\mu_{\mathbb{K}} : X \rightarrow (\text{Lie } \mathbb{K})^*$ followed by the map $\pi : (\text{Lie } \mathbb{K})^* \rightarrow (\text{Lie } K)^*$ dual of the map $\text{Lie } K \rightarrow \text{Lie } \mathbb{K}$ induced by h . The corresponding Duistermaat-Heckman measure $d\sigma_{DH}^K$ coincides, up to a constant, with the pushforward measure of the linear map of polytopes $\pi : \mu_{\mathbb{K}}(X) \rightarrow \mu_K(X)$. It is piecewise polynomial of degree $\dim X - \dim h(T)$.

In our case, the limit of the partial flag variety G/P_λ has two torus actions: by $\mathbb{T}$ and by the maximal torus T of G , related by the homomorphism

$$h : T \rightarrow \mathbb{T}, \quad t \mapsto (\alpha_{i_1}(t), \dots, \alpha_{i_N}(t))$$

with kernel being the center of G . The corresponding moment polytopes are $Q_{\underline{w}_0}(\lambda)$ and $\text{Conv}(W\lambda^*)$.

Lemma 5.12. *The pushforward of the Euclidean measure under the projection $\pi_\lambda : Q_{\underline{w}_0}(\lambda) \rightarrow \text{Conv}(W\lambda^*)$ does not depend on $\underline{w}_0$.*

Proof. Indeed, let $(\mathcal{X}, \mathcal{O}_{\mathcal{X}}(1)) \rightarrow \mathbb{A}^1$ be a one-parameter degeneration with special fiber the toric variety corresponding to the polytope $Q_{\underline{w}_0}(\lambda)$. The Duistermaat-Heckman measure depends continuously on $t \in \mathbb{C}$ and is constant for $t \neq 0$. Therefore, the measure in the limit coincides with the measure $d\sigma_{DH}$ for the K -action on G/P_λ (regarded as the coadjoint orbit of λ under a maximal compact subgroup of G).

Alternatively, one may observe that the density of the pushforward of the Euclidean measure under π_λ assigns to any $\mu \in \text{Conv}(W\lambda^*)$ the volume of its fiber

$\pi_\lambda^{-1}(\mu)$. For rational μ , this fiber is a rational convex polytope with Ehrhart function

$$n \mapsto \#(n\pi_\lambda^{-1}(\mu) \cap \mathbb{Z}^N) = \#(\pi_{n\lambda}^{-1}(n\mu) \cap \mathbb{Z}^N) = \dim V(n\lambda^*)_{n\mu},$$

the multiplicity of the weight $n\mu$ in $V(n\lambda^*)$. Thus, the density function is independent of $\underline{w}_0$. $\square$

We note that this measure contains only partial information about the polytope $Q_{\underline{w}_0}(\lambda)$. For example, the measure on Fig.1 is S_3 -symmetric. However, the corresponding Gelfand-Tsetlin polytope is not S_3 -symmetric, and only one of the special interior points on the picture is the image of a vertex of $Q_{\underline{w}_0}(\lambda)$.

For any $\mathbb{Q}$ -polarized spherical variety (X, L) , the same proof shows that the pushforward measure under the projection $\pi : Q_{\underline{w}_0}(X, L) \rightarrow \text{Conv}(W \backslash P(X, L))$ does not depend on $\underline{w}_0$.

6. CONNECTION WITH THE “UNIVERSAL” COMPACTIFIED MODULI SPACE

The Minimal Model Program, which is fully established in dimension ≤ 3 and is still conjectural in dimensions ≥ 4 , implies the existence of a complete moduli space $\overline{M}$ of stable pairs (X, D) , generalizing the Deligne-Mumford moduli space $\overline{M}_g$ of stable curves. These pairs consist of a projective variety X and a $\mathbb{Q}$ -Weil divisor D that satisfy two basic conditions:

- (i) (on singularities) (X, D) should have semi-log canonical singularities, and
- (ii) (numerical) the log canonical divisor $K_X + D$, properly defined if X is non-normal, should be ample.

In particular, every one-parameter family (X_t, D_t) of stable pairs should have, possibly after a finite base change, a unique stable limit (X_0, D_0) . See [KSB88, Ale96a, Ale96b] for more of this story.

In the situations where the varieties have an additional structure, such as a group action, the existence of such moduli space can be established in higher dimensions without using the methods of the Minimal Model Program. Two examples are pairs with toric or semiabelian group action [Ale02] and stable reductive varieties [AB02, AB03].

Below, we show that in the cases when the polytope $Q_{\underline{w}_0}(X, L)$ is integral the toric limits we have considered in this paper can be interpreted as boundary points on the compactified moduli space $\overline{M}$ of stable pairs, if it exists. First, recall the basic

Definition 6.1. Let X be a normal variety and $D = \sum a_j D_j$ be a $\mathbb{Q}$ -Weil divisor such that $0 \leq a_j \leq 1$ for all j . The pair (X, D) is canonical if some positive multiple of $K_X + D$ is an integral Cartier divisor, and if for every resolution of singularities $f : Y \rightarrow X$ with exceptional divisors F_i , we have

$$K_Y + f_{\text{strict}}^{-1} D = f^*(K_X + D) + \sum b_i F_i$$

where $b_i \geq -1$ for all i . This formula makes sense since Cartier divisors can be pulled back; the coefficients b_i are called the discrepancies.

The notion of semi-log canonical pair is a generalization of log canonical to the case when X is non-normal. We will not need this generalization since all our varieties are normal.

Let $X = G/B$ be a flag variety, E be the union of all the Schubert varieties considered as a reduced divisor, and $E^- = w_0 E$ be the opposite divisor. For a regular dominant weight λ , define the divisor $D = D_\lambda$ on X as follows:

$$D = (s), \quad \text{where } s = \sum b_{\lambda, \varphi} \in H^0(G/B, L_\lambda)$$

with the latter sum going over all elements of the dual canonical basis of $V(\lambda^*)$.

Theorem 6.2. (i) *The pair $(X, E + E^-)$ is log canonical and $K_X + E + E^-$ is rationally equivalent to 0.*
(ii) *For $0 < \varepsilon \ll 1$, the pair $(X, E + E^- + \varepsilon D)$ is stable, i.e. $(X, E + E^- + \varepsilon D)$ is log canonical and $K_X + E + E^- + \varepsilon D$ is ample.*

Proof. (i) is proved in [AB03, Thm.5.10]. Since D is ample then the divisor $K_X + E + E^- + \varepsilon D$ obviously is ample. To prove the statement about log canonical singularities, we have to prove that D does not contain any center of log canonical singularities of $E + E^-$, i.e. any image of an exceptional divisor F_i on a resolution $f : Y \rightarrow X$ that has discrepancy -1 . But such a center must be T -invariant, and s does not vanish at any T -fixed point of X . $\square$

Now, pick any family of integers (n_φ) . Then the section

$$s_t = \sum t^{n_\varphi} b_{\lambda, \varphi} \in H^0(G/B, L_\lambda)$$

defines a one-parameter family of stable pairs $(X, E + E^- + \varepsilon D_t)$. The Minimal Model Program in dimension $\dim(X) + 1$ implies that this family has a unique limit as a stable pair $(X_0, E_0 + E_0^- + \varepsilon D_0)$. It is obtained as the special fiber of the log canonical model $\mathcal{X}_{\text{can}}$ of the semistable model of the degenerating family $\mathcal{X}$.

On the other hand, the filtration $(R_{\leq m})$ of the algebra $R = R(G/B, L_\lambda)$, constructed in Proposition 2.2, can be interpreted as the data for a one parameter degeneration, in the usual way, by setting

$$n_\varphi = \min\{m \mid b_{\lambda, \varphi} \in R_{\leq m}\}.$$

Now assume that the polytope $Q_{\underline{w}_0}(\lambda)$ is integral. On the limit toric variety X_0 , the limit of $E + E^-$ is the boundary divisor ∂X_0 , i.e., the complement of the open $\mathbb{T}$ -orbit.

Indeed, on $X = G/B$ the divisor $E + E^-$ is the divisor of zeros of the product of the highest and the lowest weight vectors in $H^0(G/B, L_\lambda)$ (since λ is regular). Thus, $E_0 + E_0^-$ is the divisor of zeroes of the product of the limits of these vectors on X_0 . These limits are the sections associated with the highest (resp. lowest) weight vertices of the string polytope. The product of these sections vanishes everywhere on the boundary of X_0 , since any codimension-1 face of the string polytope contains one of these vertices, as is clear from the defining inequalities in Theorem 1.1.

This proves that $\text{supp}(E_0 + E_0^-) = \text{supp } \partial X_0$. As on any toric variety, one has $K_{X_0} + \partial X_0 \sim 0$. Applying the argument of Proposition 2.4, we see that $K_{\mathcal{X}} + \mathcal{E} + \mathcal{E}^- \sim 0$ on the family $\mathcal{X}$. Therefore, $E_0 + E_0^- \sim \partial X_0$.

This implies that the pair $(X_0, E_0 + E_0^-)$ has log canonical singularities and that its centers of log canonical singularities are precisely the closures of $\mathbb{T}$ -orbits. The limit of s_t is the section $s_0 = \sum b_{\lambda, \varphi}$, with $b_{\lambda, \varphi}$ going over the basis of $\mathbb{T}$ -eigenfunctions of $H^0(X_0, L_0)$. Again, the divisor of this section does not contain any $\mathbb{T}$ -fixed points. Hence, the pair $(X_0, E_0 + E_0^- + \varepsilon D_0)$ is stable, as predicted.

This argument can be repeated for an arbitrary polarized projective spherical variety (X, L) , provided the polytope $Q_{\underline{w}_0}(X, L)$ is integral, with s being replaced by the sum of basis vectors $b_{\lambda, \varphi}$ with λ going over the weights appearing in $H^0(X, L)$. This concludes our speculation about the connection with the “universal” moduli space of stable pairs.

Finally, we want to remark on the construction of Grossberg and Karshon [GK94] of toric limits of Bott-Samelson varieties. The latter appear naturally as resolutions of singularities of the pair $(G/B, E)$, where E , as above, is the union of Schubert varieties, considered as a reduced divisor. Every reduced decomposition $\underline{w}_0$ provides a Bott-Samelson variety $Y = Y_{\underline{w}_0}$ together with a birational morphism $f = f_{\underline{w}_0} : Y \rightarrow G/B$.

The Bott-Samelson varieties degenerate to toric limits for the torus $\mathbb{T}$, that are nonsingular toric varieties called Bott towers. In [GK94] they are described by a combinatorial structure called *twisted cube* which is not a polytope but a kind of “virtual” polytope. Translating this back to geometry, this means that the limit on Y_0 of the globally generated line bundle $f^*(L_\lambda)$ is a sheaf which is no longer globally generated. Hence, the relationship between G/B and Y gets broken in the limit, and there is only a rational map between the special fibers of degenerations.

REFERENCES

- [AB02] V. Alexeev and M. Brion, *Stable reductive varieties: affine varieties*, Invent. Math., to appear (2002), [arXiv: math.AG 0207272](#).
- [AB03] ———, *Stable reductive varieties: projective case*, Advances of Math., to appear (2003), [arXiv: math.AG 0207274](#).
- [AB04] ———, *Moduli of affine schemes with reductive group action*, J. Algebraic Geom., to appear (2004), [arXiv: math.AG 0301288](#).
- [Ale96a] V. Alexeev, *Log canonical singularities and complete moduli of stable pairs*, Preprint (1996), [alg-geom/9608013](#).
- [Ale96b] V. Alexeev, *Moduli spaces $M_{g,n}(W)$ for surfaces*, Higher-dimensional complex varieties (Trento, 1994), de Gruyter, Berlin, 1996, pp. 1–22.
- [Ale02] ———, *Complete moduli in the presence of semiabelian group action*, Ann. of Math. (2) **155** (2002), no. 3, 611–708.
- [Bat94] V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*, J. Algebraic Geom. **3** (1994), no. 3, 493–535.
- [BCFKvS00] V.V. Batyrev, I. Ciocan-Fontanine, B. Kim, and D. van Straten, *Mirror symmetry and toric degenerations of partial flag manifolds*, Acta Math. **184** (2000), no. 1, 1–39.
- [Bou87] J-F. Boutot, *Singularités rationnelles et quotients par les groupes réductifs*, Invent. Math. **88** (1987), no. 1, 65–68.
- [BZ88] A. D. Berenstein and A. V. Zelevinsky, *Tensor product multiplicities and convex polytopes in partition space*, J. Geom. Phys. **5** (1988), no. 3, 453–472.
- [BZ01] A. Berenstein and A. Zelevinsky, *Tensor product multiplicities, canonical bases and totally positive varieties*, Invent. Math. **143** (2001), no. 1, 77–128.
- [Cal02] P. Caldero, *Toric degenerations of Schubert varieties*, Transform. Groups **7** (2002), no. 1, 51–60.
- [Chi00] R. Chirivì, *LS algebras and application to Schubert varieties*, Transform. Groups **5** (2000), no. 3, 245–264.
- [CL02] P. Caldero and P. Littlemann, *Adapted algebras and standard monomials*, 2002, [arXiv:math.RT/0201026](#).
- [Dem88] M. Demazure, *Anneaux gradués normaux*, Introduction à la théorie des singularités, II, Hermann, Paris, 1988, pp. 35–68.
- [Elk78] R. Elkik, *Singularités rationnelles et déformations*, Invent. Math. **47** (1978), no. 2, 139–147.

- [Giv97] A. Givental, *Stationary phase integrals, quantum Toda lattices, flag manifolds and the mirror conjecture*, Topics in singularity theory, Amer. Math. Soc. Transl. Ser. 2, vol. 180, Amer. Math. Soc., Providence, RI, 1997, pp. 103–115.
- [GK94] M. Grossberg and Y. Karshon, *Bott towers, complete integrability, and the extended character of representations*, Duke Math. J. **76** (1994), no. 1, 23–58.
- [GL96] N. Gonciulea and V. Lakshmibai, *Degenerations of flag and Schubert varieties to toric varieties*, Transform. Groups **1** (1996), no. 3, 215–248.
- [Gro97] F.D. Grosshans, *Algebraic homogeneous spaces and invariant theory*, Springer Lecture Notes, vol. 1673, Springer, Berlin, 1997.
- [GŽ96] M.D. Grossberg and M. Žabčič, *Bott-Samelson geometry of Gelfand-Cetlin bases*, University of Georgia Math. Preprint Series **4** (1996), no. 16, 1–50.
- [Kav03] K. Kaveh, *SAGBI bases and Degeneration of Spherical Varieties to Toric Varieties*, 2003, arXiv:math.AG/0309413.
- [KM03] M. Kogan and E. Miller, *Toric degeneration of Schubert varieties and Gel'fand-Cetlin polytopes*, 2003, arXiv:math.AG/0303208.
- [KSB88] J. Kollár and N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. **91** (1988), no. 2, 299–338.
- [Lit98] P. Littelmann, *Cones, crystals, and patterns*, Transform. Groups **3** (1998), no. 2, 145–179.
- [LS86] V. Lakshmibai and C.S. Seshadri, *Geometry of G/P - V* , Journal of Algebra **100** (1986), no. 2, 462–557.
- [Oko97] V.L. Okounkov, *A remark on the Hilbert polynomial of a spherical variety*, Funct. Anal. Appl. **31** (1997), no. 2, 138–140.
- [Pop87] V.L. Popov, *Contraction of the actions of reductive algebraic groups*, Math. USSR-Sb. **58** (1987), no. 2, 311–335.
- [Stu96] B. Sturmfels, *Gröbner bases and convex polytopes*, University Lecture Series, vol. 8, American Mathematical Society, Providence, RI, 1996.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GA 30602, USA
E-mail address: `valery@math.uga.edu`

INSTITUT FOURIER, B. P. 74, 38402 SAINT-MARTIN D'HÈRES CEDEX, FRANCE
E-mail address: `Michel.Brion@ujf-grenoble.fr`